

# A Hausdorff-Measure Boundary Element Method for Scattering by Fractal Screens II: Numerical Quadrature

Andrew Gibbs<sup>1,\*</sup>, David Hewett<sup>1</sup>, Andrea Moiola<sup>2</sup>

<sup>1</sup>Department of Mathematics, University College London, UK

<sup>2</sup>Dipartimento di Matematica, Università Degli Studi di Pavia, Italy

\*Email: andrew.gibbs@ucl.ac.uk

## Abstract

In a related talk [1], the Boundary Element Method (BEM) is generalised to the case of scattering by fractal obstacles. Implementation requires evaluating integrals of singular Green's kernels over fractal domains, with respect to *Hausdorff measure*. This motivated the development of new quadrature rules, which are discussed here.

**Keywords:** Quadrature, BEM, Fractals

## 1 Introduction

We will study numerical quadrature rules for the evaluation of integrals of the form

$$I_{\Gamma, \Gamma'}[\Phi] := \int_{\Gamma} \int_{\Gamma'} \Phi(x, y) d\mathcal{H}^{d'}(y) d\mathcal{H}^d(x), \quad (1)$$

where  $\Gamma$  and  $\Gamma'$  are compact subsets of  $\mathbb{R}^2$  of Hausdorff dimension  $d > 0$  and  $d' > 0$  respectively,  $\mathcal{H}^d$  and  $\mathcal{H}^{d'}$  are the corresponding Hausdorff measures, and  $\Phi(x, y) = \frac{e^{ik|x-y|}}{4\pi|x-y|}$  is the fundamental solution for the Helmholtz equation with wavenumber  $k > 0$  in  $\mathbb{R}^3$ . (In what follows, similar results hold for the analogous problem posed in  $\mathbb{R}^2$ .)

Our motivation for approximating (1) is the *Hausdorff BEM*, which is introduced and analysed in the talk [1]. Such BEMs can model scattering by planar screens with non-integer (fractal) dimension, i.e.  $d \in (1, 2)$ .

## 2 Attractors of Iterated Function Systems

Now we describe in detail the class of fractal scatterers that we consider. An iterated function system (IFS) is a set of  $2 \leq M \in \mathbb{N}$  contracting similarities  $s_m(x) = \rho_m A_m x + \delta_m$ , with contraction factors  $\rho_m \in (0, 1)$ , rotation matrices  $A_m \in \mathbb{R}^{n \times n}$  and translations  $\delta_m \in \mathbb{R}^n$ , for  $m = 1, \dots, M$ . Saying that  $\Gamma$  is the *attractor* of the IFS means that  $\Gamma$  is the unique non-empty compact set satisfying  $\Gamma = s(\Gamma)$ , where  $s(E) := \bigcup_{m=1}^M s_m(E)$ ,  $E \subset \mathbb{R}^n$ .

Our quadrature rules are based on splitting  $\Gamma$  into sub-components, using the IFS structure.




Figure 1: Vector indices on Cantor Dust.

To describe these sub-components we adopt vector index notation. For  $\ell \in \mathbb{N}$  let  $I_\ell := \{1, \dots, M\}^\ell$ . Then for  $E \subset \mathbb{R}^n$  let  $E_0 := E$ , and for  $\mathbf{m} = (m_1, \dots, m_\ell) \in I_\ell$  define  $E_{\mathbf{m}} := s_{\mathbf{m}}(E)$  and  $s_{\mathbf{m}} := s_{m_1} \circ \dots \circ s_{m_\ell}$ . For an illustration of this notation in the case of the middle-third Cantor dust see Figure 1. We say  $\Gamma$  is *hull-disjoint* if

$$\mathcal{R} := \min_{m \neq m'} \{\text{dist}(\text{Hull}(\Gamma_m), \text{Hull}(\Gamma_{m'}))\} > 0.$$

A key ingredient is the set of vector indices

$$L_h(\Gamma) := \{\mathbf{m} = (m_1, \dots, m_\ell) \in \bigcup_{\ell \in \mathbb{N}} I_\ell : \text{diam}(\Gamma_{\mathbf{m}}) \leq h \text{ and } \text{diam}(\Gamma_{(m_1, \dots, m_{\ell-1})}) > h\}.$$

Heuristically, these indices correspond to a partition of  $\Gamma$ , where we have subdivided *just enough* so that all components have diameter no more than  $h$ . This is depicted in Figure 2.

## 3 The barycentre rule

We define the *barycentre rule* for double integrals:

$$Q_{\Gamma, \Gamma'}^h[f] := \sum_{\mathbf{m} \in L_h(\Gamma)} \sum_{\mathbf{m}' \in L_h(\Gamma')} w_{\mathbf{m}} w'_{\mathbf{m}'} f(x_{\mathbf{m}}, x'_{\mathbf{m}'}), \quad (2)$$




Figure 2: Partitioning Koch snowflake by  $L_{0,3}(\Gamma)$ . Barcentres  $x_{\mathbf{m}}$  are represented by  $\times$ .

where the weights and nodes are given by  $w_{\mathbf{m}} := \mathcal{H}^d(\Gamma_{\mathbf{m}})$  and  $x_{\mathbf{m}} := \int_{\Gamma_{\mathbf{m}}} x d\mathcal{H}^d(x) / \mathcal{H}^d(\Gamma_{\mathbf{m}})$  for  $\mathbf{m} \in L_h(\Gamma)$ , with analogous definitions for  $\Gamma'$ . The weights and nodes can be easily computed in terms of the IFS parameters, see [2, (27-29)]. For the single integral version of (2), see [2, §3.1]. In all estimates that follow,  $C$  denotes a constant which depends only on  $\Gamma$ .

**Theorem 1 (Lipschitz integrands)** [2, Theorem 3.7] *If  $L_0[f]$  and  $L_1[f]$  are the Lipschitz constants of  $f$  and  $\nabla f$  respectively in  $\text{Hull}(\Gamma) \times \text{Hull}(\Gamma')$ ,*

$$|I_{\Gamma, \Gamma'}[f] - Q_{\Gamma, \Gamma'}^h[f]| \leq CL_p[f]h^{p+1} \text{ for } p \in \{0, 1\}.$$

A result for non-diagonal entries of Hausdorff BEM matrices follows immediately:

**Corollary 2 (Smooth Galerkin integrals)** [2, Proposition 5.2]

*If  $R := \text{dist}(\text{Hull}(\Gamma), \text{Hull}(\Gamma')) > 0$ , then*

$$|I_{\Gamma, \Gamma'}[\Phi] - Q_{\Gamma, \Gamma'}^h[\Phi]| \leq Ch^2 \frac{1 + (kR)^{n/2+1}}{R^{n+1}}.$$

#### 4 Singular integrals of Laplace kernels

In Hausdorff BEM, the diagonal matrix elements correspond to (1) with  $\Gamma = \Gamma'$ . Because  $|\Phi(x, y)| \rightarrow \infty$  as  $|x - y| \rightarrow 0$ , the rule (2) cannot be directly applied to (1) in this case. We will derive a new method for evaluating the singular (Laplace) component of (2), denoted  $\Phi_0(x, y) := |x - y|^{-1}$ . Then, to evaluate (1) with  $\Gamma = \Gamma'$ , we use a *singularity subtraction*

technique, by considering the Lipschitz continuous function  $\Phi_* := \Phi - \Phi_0$ , and splitting the integral as follows

$$I_{\Gamma, \Gamma}[\Phi] = I_{\Gamma, \Gamma}[\Phi_0] + I_{\Gamma, \Gamma}[\Phi_*], \quad (3)$$

and evaluating both components separately.

By exploiting the self-similarity of  $\Gamma$ , we can express  $I_{\Gamma, \Gamma}[\Phi_0]$  as a linear function of  $I_{\Gamma_m, \Gamma_m}[\Phi_0]$  for  $m = 1, \dots, M$ , which leads to

$$I_{\Gamma, \Gamma}[\Phi_0] = \frac{\sum_{m=1}^M \sum_{m' \neq m}^M I_{\Gamma_m, \Gamma_{m'}}[\Phi_0]}{1 - \sum_{m=1}^M \rho_m^{2d-1}}, \quad (4)$$

representing a singular integral as a linear combination of smooth integrals. The smooth integrals of (4) can be approximated using (2); we denote this approximation by  $Q_{\Gamma, \Gamma, 0}^h$ .

**Theorem 3 (Singular Laplace-type integrals)**

[2, Corollary 4.7] *If  $\Gamma$  is Hull-disjoint, then*

$$|I_{\Gamma, \Gamma}[\Phi_0] - Q_{\Gamma, \Gamma, 0}^h| \leq Ch^2 \mathcal{R}^{-3} \left( 1 - \sum_{m=1}^M \rho_m^{2d-1} \right)^{-1}.$$

#### 5 Approximating (1)

Noting the decomposition (3), Theorem 3 states that  $I_{\Gamma, \Gamma}[\Phi_0]$  can be estimated with  $O(h^2)$  error, provided  $\Gamma$  is hull-disjoint.

Since  $\Phi_* \in C^{0,1}(\mathbb{R}^n) \setminus C^{1,1}(\mathbb{R}^n)$ , Theorem 1 suggests  $|I_{\Gamma, \Gamma}[\Phi_*] - Q_{\Gamma, \Gamma}^h[\Phi_*]| = O(h)$ . With further work it can be shown that this is actually  $O(h^2)$ , when (i)  $\Gamma$  is hull-disjoint and (ii)  $\rho_1 = \dots = \rho_M$ . Hence using (3) we can approximate  $I_{\Gamma, \Gamma}[\Phi]$  with  $O(h^2)$  accuracy (see [2, §5] for details).

Furthermore, numerical experiments [2, §6] suggest  $O(h^2)$  convergence for fractals which violate either or both of the conditions (i)–(ii).

#### References

- [1] S. N. Chandler-Wilde, A. Caetano, A. Gibbs, D. P. Hewett and A. Moiola, A Hausdorff-Measure Boundary Element Method for Scattering by Fractal Screens I: Numerical Analysis, in *Proceedings of the 15th International Conference on Mathematical and Numerical Aspects of Wave Propagation, Paris, France, 25-29 July 2022*.
- [2] A. Gibbs, D. P. Hewett and A. Moiola, Numerical quadrature for singular integrals on fractals, *arXiv* **2112.11793** (2021).